

Iodine Tests

lodine deficiency is re-emerging in many locations around the world. Since this essential mineral is required for the synthesis of thyroid hormones, the major indication of an iodine deficiency include hypothyroid symptoms. Furthermore, an iodine deficiency during pregnancy can result in cretism, lowered intellectual quotients (IQ), mental retardation, autism, increased infant mortality, growth stunting and impaired movement. Iodine may also play a role in the maintenance of healthy breast tissue and cancer prevention. Its antioxidant, immune regulating and estrogen modulating functions may be mechanisms by which these properties can be explained. Its antioxidant, immune regulating and estrogen modulating functioners are aware of the effects of an iodine deficiency and investigate it in their patients.

Table 1: Symptoms and Conditions Associated with an Iodine Deficiency	
Hypothyroidism	Decreased fertility
Goiter	Prostate, endometrial, ovarian, and breast cancers
Fibrocystic breast disease	Cardiovascular disease
Estrogen imbalance	Cretism
Fatigue	Lowered intellectual quotients (IQ)
Depression	Increased perinatal death & infant mortality
Dry skin	Growth stunting
Weight gain	Impaired movement
Myalgia	Speech and hearing problems

Iodine Deficiency

Some believe that iodine deficiency is a thing of the past, that it was resolved in the 1930s with the iodisation practices. In actual fact current public health recommendations to restrict salt intake, combined with a decline in the use of iodophors by the dairy industry and in commercial bread production, has actually lead to the re-emergence of this mineral deficiency in many locations around the world. For example, recent research has indicated that severe iodine deficiency has more than quadrupled in America over the last 20 years; increasing from 2.6% to 11.7%. Even more disconcerting is the observation that nearly half of the pregnant women in these areas also have this mineral deficiency. Attention to this matter by practitioner is therefore urgently required because of the serious implications for maternal and child health.

Iodine Tests Available

- **Iodine, random (Urinary):** Measures iodine levels from a urinary (spot or 24 hour) specimen.
- **Iodine Loading Test (Urinary):** Measures both a) iodine levels from a urinary spot specimen and b) the percentage of iodine (% excretion) that is excreted over a 24 hour time period, after the consumption of a large loading dose of iodine (50mg).

Which Test Should I Choose?

The Random Iodine Test: This test is the standard procedure used to detect iodine deficiencies. It measures the iodine levels in a random urinary specimen. The disadvantage of this diagnostic procedure however, is that it is not a sensitive test for detecting mild or moderate iodine deficiencies.

The lodine Loading Test: This test assesses iodine deficiency using two different methods. Firstly, it measures iodine levels in a morning spot urinary specimen (i.e. equivalent to the random iodine test). In addition, the loading test analyses iodine deficiency using a much more sensitive technique. For this part of the test, 50mg of an iodine/iodide mixture is given as a loading dose and the amount of iodine excreted in the urine over the next 24 hours is measured. If the patient is iodine deficient, the body holds on to the iodine and only a small quantity of the mineral is excreted into the urine (i.e. giving a low % excretion). In contrast, if the patient has sufficient iodine levels, the body does not retain the iodine, and therefore the majority of the iodine dosage is excreted into the 24 hour urinary sample (i.e. an adequate % excretion). In an iodine sufficient state, approximately 90% of the 50mg dose of iodine/iodide would be excreted (i.e. 45mg) and 10% of the iodine would be retained (i.e. 5mg). This test was invented by Dr Brownstein, author of the book *lodine: Why You Need It, Why You Can't Live Without It*. In contrast to the random iodine test, the iodine loading test is able to detect mild and moderate, as well as severe deficiencies.

How to order a test kit:

Phone Customer Service on 1300 688 522.

Related Tests

- Mineral and Metal Screens
- Thyroid Hormone Tests

Technical support

NutriPATH offers technical assistance by experienced clinical consultants.



Phone 1300 688 522 for further details www.nutripath.com.au