

P: 1300 688 522 E: info@nutripath.com.au

## Dr.SAMPLE REPORT TEST HEALTH CENTRE 123 TEST STREET BURWOOD VIC 3125

## SAMPLE REPORT 09-May-1990 Female

16 HARKER STREET BURWOOD VIC 3125

LAB ID : UR NO. : 3814123

Collection Date : 09-May-2022 Received Date:09-May-2022



BIOCHEMISTRY BLOOD - SERUM Result Range Units LIVER FUNCTION TESTS umol/L **BILIRUBIN (TOTAL)** 3 - 15 6 35 - 110 units/I ALP 49 units/L 41 \*\*H 5 - 35 GGT units/L ALT 5 - 30 22 AST 32 10 - 35 units/L **PROTEIN - TOTAL** 78 60 - 83 g/L **ALBUMIN** 44 35 - 50 g/L **GLOBULIN** g/L 34

GGT elevation is sensitive, but non-specific, for hepatobilary diseases. Biliary obstruction can be intrahepatic, extrahepatic or infiltrative. Microsomal GGT can also be induced by many drugs (phenytoin etc.) and ETOH, this change is reversable.

AMYLASE	1	0 - 100	units/L	•
LIPASE	42	13 - 60	units/L	

GIT ASSESSMENTS							
	Result	Range	Units				
Pancreatic Elastase 1	>500	> 200	ug/g				

## **Digestive Markers Comment**

PANCREATIC ELASTASE: Normal exocrine pancreatic function.

Pancreatic Elastase reflects trypsin, chymotrypsin, amylase and lipase activity. This test is not affected by supplements of pancreatic enzymes.

Healthy individuals produce on average 500 ug/g of PE-1. Thus, levels below 500 ug/g and above 200 ug/g suggest a deviation from optimal pancreatic function.

The clinician should therefore consider digestive enzyme supplementation if one or more of the following conditions is present:

Loose watery stools, Undigested food in the stools, Post-prandial abdominal pain, Nausea or colicky abdominal pain, Gastroesophageal reflux symptoms, Bloating or food intolerance.

LFT Comment

GAMMA-GLUTAMYLTRANSFERASE COMMENT: