



SAMPLE REPORT

09-May-1990 Female

P: 1300 688 522
E: info@nutripath.com.au

16 HARKER STREET
BURWOOD VIC 3125

Dr.SAMPLE REPORT
TEST HEALTH CENTRE
123 TEST STREET
BURWOOD VIC 3125

LAB ID : 3814142
UR NO. :
Collection Date : 09-May-2022
Received Date:09-May-2022



3814142

IMMUNOLOGY

BLOOD - SERUM Result Range Units

CARDIOLIPIN AB

Cardiolipin IgG:	<1	GPL-U/mL	(R.R: <10)
Beta 2 Glycoprotein IgG:	<1	U/mL	(R.R: <7)

Interpretation:

Cardiolipin IgG

Negative	< 10	GPL-U/mL
Weak Positive	10 - 40	GPL-U/mL
POSITIVE	> 40	GPL-U/mL

beta 2 Glycoprotein IgG

Negative	< 7	U/mL
Weak Positive	7 - 10	U/mL
POSITIVE	> 10	U/mL

Test performed by accredited laboratory NATA: 2133

Comments:

Beta 2 glycoprotein 1 (B2GP1) IgG antibodies are more strongly associated with the presence of antiphospholipid syndrome (APS) than anti-cardiolipin antibodies. However the absence of B2GP1 IgG antibodies does not necessarily exclude APS.

Anti-cardiolipin IgG antibodies may be associated with one or more of the following: arterial and/or venous thrombosis, recurrent pregnancy loss, low platelet counts.

Some cases are associated with systemic autoimmune disease e.g. SLE.

Low level antibodies (<40 IU/mL) are generally of uncertain clinical significance and may occur transiently in some infections.

FURTHER ASSESSMENTS:

Other tests which may be useful include lupus anticoagulant (additional specimen required) and ANA/ENA/DNA.

Tests ordered: ACL,IMPEI,CFee